

Derivation of Romberg's Rule Using N-Subdivisions and Applications of Romberg's Rule

M. N. Salim 

Division of Basic Science, College of Agriculture and Forestry, University of Mosul, Mosul, Iraq

Article information

Article history:

Received: December 08, 2024

Revised: January 19, 2025

Accepted: January 26, 2025

Available online: April 01, 2025

Keywords:

Romberg Rule

Subdivisions

$O(h^2)$

Trapezoid Rule

Matrix

Correspondence:

Mustafa Nadhim Salim

mustafa.nadhim@uomosul.edu.iq

Abstract

By using a large number of subdivisions n to obtain good accuracy in approximating the integral $\int_a^b f(x) dx$ we note that an increase in the value of n does not necessarily mean an increase in accuracy in the result or even an improvement in the result. In addition, increasing the number n means an increase in the number of times the function values are calculated. The importance of Romberg's rule lies in improving the results obtained using the previous rules and obtaining high accuracy in arriving at approximate results faster and with a small amount of error. The reason for this is the high rank of the Romberg rule compared to the trapezoid rule, for example, which is not accurate in its results because the cutting error for this rule is of the order $O(h^2)$ only.

DOI: [10.33899/edusj.2025.155630.1520](https://doi.org/10.33899/edusj.2025.155630.1520), ©Authors, 2025, College of Education for Pure Science, University of Mosul.

This is an open access article under the CC BY 4.0 license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Romberg's rule is a numerical integration method that provides a more precise approximation of definite integrals by combining Richardson extrapolation with the trapezoidal rule. It is especially helpful for smooth functions that polynomials can accurately approximate.

Romberg's rule yields a more accurate estimate than the trapezoidal rule by itself when combined with Richardson extrapolation and also Efficiency: Compared to other approaches, it needs fewer function evaluations, particularly for smooth functions. The integral $\int_a^b f(x) dx$ may be approximated with excellent precision by using a large number of subdivisions n , which divide the time period into subintervals and yield accurate and correct results. Romberg's rule is important since it yields approximate findings. Compared to earlier techniques that rely on employing fewer divisions, this approach is quicker and has a lesser inaccuracy since it uses high ranks. A useful method for estimating definite integrals, Romberg's rule is a potent numerical integration approach that improves the accuracy of the trapezoidal rule by Richardson extrapolation.

In this research, the following lower triangular matrix of integers is produced using Romberg's method and is all an approximation of the result of the above integral:

$R(1,1)$

$R(2,1)$ $R(2,2)$

$R(n,1)$ $R(n,2) \dots$ $R(n,n)$

where $R(1,1)$, $R(2,1)$, and so on for each member of the matrix indicate the first, second, and subsequent approximate values, respectively, using the trapezoid rule.

2. Research Method (Bold, 12 pt)

The following relationships, which rely on the matrix's argument, can be used to locate the elements of the matrix. The trapezoid rule is used to determine the initial value.

$$R(1,1) = \frac{b-a}{2} [f(a) + f(b)]$$

The first value determines the remainder

$$R(k, 1) = \frac{1}{2} [R(k-1, 1) + h_{k-1} \sum_{i=1}^{2^{k-2}} f(a + (i-0.5)h_{k-1})]$$

$$h_{k-1} = \frac{b-a}{2^{k-2}}, \quad k = 2, 3, \dots, n$$

The values obtained from the interpolation are shown in the other columns. The remaining elements in the matrix may be found using the following law:

$$R(k, i) = \frac{4^{i-1} R(k, i-1) - R(k-1, i-1)}{4^{i-1} - 1}$$

$k, i = 2, 3, \dots, n$




Figure 1. The elements in the matrix Romberg's rule

3. Results And Discussion

3.1 Derivation of Romberg Rule

When the value of the function and its derivatives fall in the interval $[a, b]$, Romperk's method provides a multiple arithmetic approximation of the integral by bisecting h once more.

$$\int_a^b f(x) dx \cong \frac{h}{2} \left[f_0 + f_n + 2 \sum_{i=1}^{n-1} f_i \right] + \sum_{j=1}^{\infty} \alpha_j h^{2j}$$

where h is the period's duration and α_j is the set of all j values independent of h .

Assuming $T_{k,0}$, $k=0, 1, 2, \dots$ represents the use of the preceding approximation and $h_k = \frac{b-a}{2^k}$

$$I = T_{k,0} + \alpha_1 h_k^2 + \alpha_2 h_k^4 + \alpha_3 h_k^6 + \dots \quad (1)$$

The number of subintervals is now divided in half and doubled.

$$\begin{aligned} I &= T_{k+1,0} + \alpha_1 \left(\frac{h_k}{2}\right)^2 + \alpha_2 \left(\frac{h_k}{2}\right)^4 + \alpha_3 \left(\frac{h_k}{2}\right)^6 + \dots \\ &= T_{k+1,0} + \frac{1}{4} \alpha_1 h_k^2 + \frac{1}{16} \alpha_2 h_k^4 + \frac{1}{64} \alpha_3 h_k^6 + \dots \quad (2) \end{aligned}$$

We may now remove the components that include h^2 . To do this, multiply equation (2) by (4) and deduct equation (1) from it. The result is

$$\begin{aligned} 3I &= 4T_{k+1,0} - T_{k,0} - \frac{3}{4} \alpha_2 h_k^4 - \frac{15}{16} \alpha_3 h_k^6 + \dots \\ I &= \frac{1}{3} (4T_{k+1,0} - T_{k,0}) - \frac{1}{4} \alpha_2 h_k^4 - \frac{5}{16} \alpha_3 h_k^6 \dots \quad (3) \end{aligned}$$

Where the error value is represented by $\frac{1}{4} \alpha_2 h_k^4$ and

$$T_{k+1,1} = \frac{4T_{k+1,0} - T_{k,0}}{3}$$

It's called completing your first romper.

Equation (3) makes it evident that the value of I is approximated to be of the fourth order, which makes it more accurate ($T(k,0), T(k+1,0)$) than the second order. As a result, estimations of the sixth order may be obtained by repeating the external interpolation procedure.

$$I = T_{k,1} + \beta_1 h_k^4 + \beta_2 h_k^6 + \dots \dots (4)$$

Since β_1, β_2 is independent on of h, we may derive it by halving h and doubling the subintervals.

$$I = T_{k+1,1} + \beta_1 \left(\frac{h_k}{2}\right)^4 + \beta_2 \left(\frac{h_k}{2}\right)^6 + \dots$$

$$= T_{k+1,1} + \frac{1}{16} \beta_1 h_k^4 + \frac{1}{64} \beta_2 h_k^6 + \dots \dots (5)$$

After subtracting equation (4) from equation (5) and multiplying it by (4), we obtain

$$I = \frac{1}{15} (16T_{k+1,1} - T_{k,1}) - \frac{1}{20} \beta_1 h_k^6$$

The second Romberg interpolation, or $T(k+1,2)$, is the prior equation. From there, we proceed to get the best estimate.

Common law

$$I = T_{k+1,j+1} + O(h^{2j+4})$$

When $J=0$ Romperk I $T(k+1,1)+O(h^4)$

When $J=1$ Romperk II $T(k+1,2)+O(h^4)$

When $J=2$ Romperk III $T(k+1,3)+O(h^4)$

$$\text{So it is possible to calculate } T_{k+1,j+1} = \frac{4^{j+1} T_{k+1,j} - T_{k,j}}{4^{j+1} - 1}$$

By using Romberg's rule $\int_0^2 x^2 dx$, where $n=3$ we have

$$R(1,1)$$

$$R(2,1) \quad R(2,2)$$

$$R(3,1) \quad R(3,2) \quad R(3,3)$$

$$R(1,1) = \frac{b-a}{2} [F(a) + F(b)]$$

$$= \frac{2-0}{2} [0 + 4] = 4$$

$$h_{k-1} = \frac{b-a}{2^{k-2}}, \quad K=2$$

$$h_1 = \frac{b-a}{2^0} = h_1 = 2$$

$$h_2 = \frac{2}{2} = 1$$

$$R(2,1) = \frac{1}{2} [R(1,1) + h_1 \sum_{i=1}^1 f(0 + 0.5)(2)]$$

$$= \frac{1}{2} [4 + 2f(0 + 0.5)(2)] = 3$$

$$R(3,1) = \frac{1}{2} [R(2,1) + h_2 \sum_{i=1}^2 f(a + (i - 0.5)h_2)]$$

$$= \frac{1}{2} \left[3 + f(0 + 0.5) + f\left(0 + \frac{3}{2}\right) \right]$$

$$= \frac{1}{2} \left[3 + \frac{1}{4} + \frac{9}{4} \right] = 2.75$$

We apply the following formula to get the matrix's remaining members

$$R(k, i) = \frac{4^{i-1} R(k, i-1) - R(k-1, i-1)}{4^{i-1} - 1}$$

$$R(2,2) = \frac{4R(2,1) - R(1,1)}{4-1}$$

$$= \frac{4(3) - 4}{3}$$

$$= \frac{8}{3} = 2.66667$$

$$R(3,2) = \frac{4R(3,1) - R(2,1)}{4-1}$$

$$= 2.66667$$

$$R(3,3) = \frac{4^{3-1} R(3,2) - R(2,2)}{4^{3-1} - 1}$$

$$= \frac{16(2.66667) - 2.66667}{15}$$

$$= 2.66667$$

This is equivalent to the integral's actual value

$$\int_0^2 x^2 dx = \frac{x^3}{3} \Big|_0^2 \\ = \frac{8}{3} \cong 2.6666$$

If the following table is used to represent the speed of an automobile at time t, then

T	0	12	24	36	48	60	72	84	96	108	120
V	0	3.60	10.08	18.90	21.60	18.54	10.26	5.30	4.50	5.40	9.00

Determine the distance the car travels in a time limit of no more than two minutes using Romberg's rule.

$$R(1,1) = \frac{b-a}{2} [f(a) + f(b)]$$

$$f(a) = f(0) = 0$$

$$f(b) = f(120) = 120$$

$$R(1,1) = \frac{120-0}{2} [0 + 120] = 7200$$

$$R(k, 1) = \frac{1}{2} [R(k-1, 1) + h_{k-1} \sum_{i=1}^{2^{k-2}} f(a + (i-0.5))]$$

$$h_{k-1} = \frac{b-a}{2^{k-2}}, \quad k = 2, 3, \dots, n$$

$$h_{k-1} = \frac{b-a}{2^{k-2}}, \quad k = 2$$

$$h_{2-1} = \frac{120-0}{2^0} = 120$$

$$R(2,1) = \frac{1}{2} [R(1,1) + 120 \sum_{i=1}^{2^0} f(0 + (i-0.5))]$$

$$= \frac{1}{2} [7200 + 120(0.5)] = 216000$$

$$R(k, i) = \frac{4^{i-1} R(k, i-1) - R(k-1, i-1)}{4^{i-1} - 1}$$

$$R(2,2) = \frac{4^{2-1} R(2,1) - R(1,1)}{4^{2-1} - 1}$$

$$= \frac{4(216000) - 7200}{3} \approx 280600$$

The results are in the form of the following matrix:

7200	0
216000	280600

Table 1. Romberg's rule is better to the trapezoidal rule because it allows for faster convergence to the integer value.

	O(h)	h²	h⁴	h⁶	h⁸
I					
h		T _{0,0}			
$\frac{h}{2}$		T _{0,1}	T _{1,0}		
$\frac{h}{4}$		T _{0,2}	T _{1,1}	T _{2,0}	
$\frac{h}{8}$		T _{0,3}	T _{1,2}	T _{2,1}	T_{3,0}

3.2 Algorithm of Romberg Rule

Step 1: type f, a, b, and n.

Step 2: We determine the first integral's value $R(1,1) = \frac{b-a}{2} [F(A) + F(B)]$

Step 3: The values of h are determined utilizing $h = \frac{b-a}{(2^{k-2})}$

Step 4: For any value of k and i, we compute the following integral's value.

$$R(k, 1) = \frac{1}{2} [R(k-1, 1) + h_{k-1} \sum_{i=1}^{2^{k-2}} f(a + (i-0.5))]$$

Where $k = 2, 3, \dots, n$, $i = 1, 2, \dots, 2^{k-2}$

Step 5: We calculate Romperk's rule

$$R(k, i) = \frac{4^{i-1} R(k, i-1) - R(k-1, i-1)}{4^{i-1} - 1}$$

K, $i=2,3,\dots,n$

Step 6: Print the integral value

3.3 Program of Romberg Rule

```
f= input('f='); %function needed to integrate
a= input('a='); % start point of interval
b= input('b='); % end point of interval
n= input('n=');
r(1,1)=((b-a)/2)*(subs(f,a)+subs(f,b));
for k=2:n
    h=(b-a)/(2^(k-2));
    s=0;
    for i=1:2^(k-2)
        s=s+ subs(f,(a+(i-0.5)*h));
    end
    r(k,1)=0.5*(r(k-1,1)+h*s);
end
for i=2:n
    for k=i:n
        r(k,i)=((4^(i-1)*r(k, i-1) - r(k-1, i-1))/(4^(i-1)-1));
    end
end
r
```

4. Conclusion

The use of a high number of subdivisions, n , in the approximation of the integral is one of the most significant characteristics of using $\int_a^b f(x) dx$. Romberg's rule has a significant effect on the outcomes as higher n increases the number of times the function values are computed. As a consequence, it may be applied to mathematical analyses of a variety of mathematical issues and produces fast, highly accurate answers with minimal mistakes, improving the outcomes of the rules before.

5. Acknowledgements

The researchers extend their thanks and gratitude to the Presidency of the University of Mosul and the College of Agriculture & Forestry for facilitating the task of completing the research and conducting laboratory tests.

6. References

- [1] M. Bohner and A. Peterson, *Dynamic equations on time scales: An introduction with applications*. Springer Science & Business Media, 2001.
- [2] Khasnis, A., and R. M. Gokula. "Romberg's Test." *Journal of postgraduate medicine* 49.2 (2003): 169-172.
- [3] K. Diethelm, N. J. Ford, and A. D. Freed, "Detailed error analysis for a fractional Adams method," *Numerical Algorithms*, vol. 36, no. 1. pp. 31–52, 2004. doi: 10.1023/B:NUMA.0000027736.85078.be
- [4] S. Momani and N. Shawagfeh, "Decomposition method for solving fractional Riccati differential equations," *Applied Mathematics and Computation*, vol. 182, no. 2. pp. 1083–1092, 2006. doi: 10.1016/j.amc.2006.05.008.
- [5] Tseng, C. C., & Lee, S. L. (2008). Digital IIR integrator design using recursive Romberg integration rule and fractional sample delay. *Signal Processing*, 88(9), 2222-2233.
- [6] Tseng, Chien-Cheng, and Su-Ling Lee. "Digital IIR integrator design using recursive Romberg integration rule and fractional sample delay." *Signal Processing* 88.9 (2008): 2222-2233.
- [7] Polla, Gerardus. "Comparison of Approximation Accuracy and time Integral Process between Simpson Adaptive Method and Romberg Method." *Journal of Interdisciplinary Mathematics* 18.5 (2015): 459-470.
- [8] D. Pachpatte, "On some dynamic inequalities in two variables on time scales," *AIP Conf. Proc.*, vol. 2061, no. January, 2019, doi: 10.1063/1.5086626
- [9] S. Kumar et al., "An efficient numerical method for fractional SIR epidemic model of infectious disease by using Bernstein wavelets," *Mathematics*, vol. 8, no. 4, p. 558, 2020.

اشتقاق قاعدة رومبرك باستخدام التقسيمات الجزئية n وتطبيقات قاعدة رومبرك الرياضية

مصطفى ناظم سالم

فرع العلوم الاساسية، كلية الزراعة والغابات، جامعة الموصل، الموصل، العراق

المستخلص:

باستخدام عدد كبير من التقسيمات الجزئية n لغرض الحصول على دقة جيدة في التقريب للتكامل $\int_a^b f(x) dx$ نلاحظ بأنه ليس بالضرورة أن يعني الزيادة في قيمة n زيادة في الدقة في النتيجة أو حتى تحسين في النتيجة إضافة إلى ذلك إن زيادة العدد n تعني زيادة بعدد المرات التي تحسب فيها قيم الدالة، إن أهمية قاعدة رومبرك تكمن في تحسين النتائج التي تم الحصول عليها باستخدام القواعد السابقة والحصول على دقة عالية في الوصول إلى نتائج تقريبية بشكل أسرع وبمقدار خطأ بسيط، والسبب في ذلك هي الرتبة العالية لقاعدة رومبرك بالمقارنة مع قاعدة شبه المنحرف مثلاً التي لا تكون دقيقة في نتائجها وذلك لأن خطأ القطع لهذه القاعدة من الرتبة $O(h^2)$ فقط.